

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 190 205 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**(45) Date of publication of patent specification: **21.10.92** (51) Int. Cl.<sup>5</sup>: **G01N 33/53, G01N 33/68**(21) Application number: **85903646.9**(22) Date of filing: **23.07.85**(86) International application number:  
**PCT/AU85/00165**(87) International publication number:  
**WO 86/00991 (13.02.86 86/04)****(54) METHOD FOR DETERMINING MIMOTOPES.**(30) Priority: **24.07.84 AU 6188/84**(43) Date of publication of application:  
**13.08.86 Bulletin 86/33**(45) Publication of the grant of the patent:  
**21.10.92 Bulletin 92/43**(84) Designated Contracting States:  
**AT BE CH DE FR GB IT LI LU NL SE**(56) References cited:  
**WO-A-84/02983**  
**WO-A-84/03564**  
**WO-A-85/02121**  
**AU-A- 8 425 428**  
**AU-B- 7 473 787****PROCEEDINGS OF THE NATL. ACADEMY OF  
SCIENCE USA, vol. 81, July 1984;**  
**H.M.GEYSEN et al., pp. 3998-4002****CHEMICAL ABSTRACTS, vol. 101, no. 7, 13**  
**August 1984, Columbus, OH (US);****J.WEHLAND et al., p. 455, no. 52962p****PROCEEDINGS OF THE NATL. ACADEMY OF  
SCIENCE USA, vol. 82, January 1985;**  
**H.M.GEYSEN et al., pp. 178-182****SCIENTIFIC AMERICAN, February 1983;**  
**R.A.LERNER, pp. 48-56**(73) Proprietor: **COSELCO MIMOTOPES PTY. LTD.**  
**Corner Duerdin and Martin Streets**  
**Clayton, VIC 3168(AU)**(72) Inventor: **GEYSEN, Hendrik, Mario**  
**10 Janden Close**  
**Knoxfield, VIC 3180(AU)**(74) Representative: **Allard, Susan Joyce et al**  
**BOULT, WADE & TENNANT, 27 Furnival**  
**Street**  
**London EC4A 1PQ(GB)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Rank Xerox (UK) Business Services

**EP 0 190 205 B1****BEST AVAILABLE COPY**

## Description

This invention relates to a method of detecting or determining a sequence of monomer molecules which corresponds to an epitope on an antigen, or which has the equivalent molecular topology to that epitope.

As used throughout this specification, the terms listed below have the following meanings:-

epitope:

the specific surface of an antigen molecule which is delineated by the area of interaction with an antibody molecule.

catamer:

a polymer molecule which has a precisely defined sequence and which is formed by the condensation of a precise number of small molecules. Note that this term includes molecules in which different types of condensation reactions are used to join the small molecules. A number prefixed to the word 'catamer' indicates that the catamer is formed by the condensation of the indicated number of small molecules, for example, 8-catamer means that the catamer is made up from eight small molecules. Examples of catamers include any given peptide, and any given oligo-saccharide.

monomer:

a member of the set of small molecules which can be condensed together to form a catamer. The set of monomers includes but is not restricted to, for example, the set of common L-amino acids, the set of D-amino acids, the set of synthetic amino acids, the set of nucleotides, and the set of pentoses and hexoses.

peptide:

a catamer in which the small molecules are alpha-amino acids which are joined together through peptide bonds. In the context of this specification it should be appreciated that the amino acids may be the L-optical isomer or the D-optical isomer.

mimotope:

catamer which in at least one of its conformations has a surface region with the equivalent molecular topology to the epitope of which it is a mimic.

complementary:

refers to the matching together of the reacting surfaces of an antibody paratope and its particular epitope. These surfaces must match during the binding of an antibody paratope with its epitope. Thus, the paratope and its epitope can be described as complementary, and furthermore, the contact surface characteristics are complementary to each other.

paratope:

the combining site of an antibody which is complementary to a particular epitope.

It is already well known that:-

1. an antigen is a macromolecule such as a protein or polysaccharide which is usually, but not necessarily, foreign to a human or animal and when introduced into the human or animal body, stimulates either a humoral or cellular immune response, or both;
2. the specific region of the antigen to which antibodies bind is known as the epitope;
3. there may be more than one epitope on an antigen molecule;
4. the overall humoral response of the body to an antigen is to produce antibodies against the various epitopes of the antigen; each of the individual antibodies is a monoclonal antibody and by definition is produced from a group of cells which come from the multiplication and differentiation of a single progenitor B-cell;
5. the paratope of an antibody molecule is defined as the molecular surface of the antibody combining site which is the contact surface between epitope and the antibody;
6. there may be produced against a given epitope of the antigen more than one antibody which differ in their paratopes; each paratope can be characterised by its specificity for the complementary epitope; and
7. antibodies produced in response to a challenge by antigen can be further characterised in that some may neutralize directly by binding with the antigen; others may form aggregates which can be dealt with by the macrophage system; and yet others may fail to exhibit any measurable biological effects.

It is a primary object of the present invention to detect or determine the one or more short sequences of monomers which equate with an epitope of an antigen, such that these sequences can combine with the same antibody as the epitope. These catamers would be the mimotopes of the epitope. This information is invaluable in the design of synthetic catamer vaccines against diseases and for the design of very specific diagnostic and therapeutic agents.

The most usual group of small molecules which may be condensed together to form a catamer is the group of alpha-amino acids. However, other molecules which are consistent with the chemistry chosen may

be used. For example, beta-amino acids may be used with advantage to add an extra -C- bond spacer at precisely controlled positions in the catamer.

PNAS, 81 (1984), p3998-4002 discloses a procedure for the rapid synthesis of hundreds of short peptides derived from the known sequence of FMDV type O<sub>1</sub>. These peptides can then be screened for immunogenicity.

EMBO journal, Vol 3, no. 6, (1984) p1295-1300 discloses the characterization of the epitope of the rat monoclonal antibody YL 1/2. This characterization is carried out using synthetic peptides spanning the carboxy-terminus of tyrosinated  $\alpha$ -tubulin.

According to the present invention there is provided a method of determining the sequence of monomers selected from a defined set of monomers which is a conformational equivalent of an epitope which is complementary to a particular paratope of an antibody of interest, the method comprising the steps of:

(i) synthesizing a plurality of catamer preparations; each of said catamer preparations consisting of a plurality of catamers made of a defined set of monomers and in which (i) the composition at two or more designated positions in each catamer is known and is constant for each catamer in the catamer preparation, and (ii) the composition at each of the remaining positions are randomly made up from the members of said defined set of monomers; and said plurality of catamer preparations comprising preparations in which the monomer(s) at the designated positions are systematically varied to contain members from said defined set of monomers;

(ii) contacting each of said plurality of catamer preparations with the antibody of interest; and

(iii) detecting the presence or absence of binding between each of said plurality of catamer preparations and said given antibody; and

(iv) determining a partial sequence of monomers for the conformational equivalent of an epitope which is complementary to the particular paratope of said antibody from the known composition of said two or more designated positions in each of said plurality of catamer preparations that bind to said antibody.

Preferably, the method also comprises the further steps of:-

(v) synthesizing a further plurality of catamer preparations as defined in claim 1 and which include the composition of the partial sequence as determined in said step (iv) as initial designated position, each of said further plurality of catamer preparations having at least one or more additional position in each catamer as a designated position in which the monomer is known; and

(vi) repeating (ii), (iii), and (iv) as defined in claim 1 in respect of said further plurality of catamer preparations.

The procedures of steps (v) and (vi) may be repeated until all positions in each catamer preparation have been designated; and all deduced mimotopes may then be synthesised to determine from their relative ability to react with the antibody of interest which are the better mimotopes for the epitope corresponding to this given antibody.

The method of the present invention is based on the realisation that a given antibody will specifically react with a catamer which is the mimotope of the epitope to which the antibody is directed. It further relies on the knowledge that modern techniques of immunology allow the detection of reactions between an antibody and its epitope when both are present in very small amounts. Binding has been detected when an antibody and its epitope are present in amounts of about 1 pmole. Thus, if an antibody is presented with a mixture of catamers, amongst which is a mimotope to its epitope then it will be seen to react with the mixture, binding specifically to the mimotope. The identity of mimotopes is deduced from the known monomer composition at the designated positions of reacting catamer preparations. A complete mimotope would be expected to be some combination of the reacting components deduced in this way.

It will be apparent that the method of this invention requires no previous information about the nature of the antigen and in particular it requires no foreknowledge of the sequence of monomers which make up the antigen. In fact, it is not necessary for the application of this invention to know the source or the identity of the antigen to which the antibody is directed. Furthermore, this invention makes no assumptions about the epitope which stimulated the production of the particular antibody. This method will identify mimotopes of discontinuous as well as continuous epitopes. Because of the very nature of the method of the invention it will be appreciated that it applies to any antigen, which in part or in total is non-proteinaceous, whose epitopes can be mimicked by a suitable catamer. Mimotopes may or may not be made from members of the same set of monomers which make up the antigen.

Preferably the method of the present invention is carried out by screening a plurality of synthesized catamer preparations against an antibody of interest. Ideally the antibody will be a monoclonal antibody which can be prepared by any of the usual methods. Polyclonal antiserum from humans and animals may be used if a source of monoclonal antibody is not available, however, analysis of the resulting data may be complicated because the reactions observed could be from more than one monoclonal antibody. When

using polyclonal antiserum it may be necessary to reduce the antibody diversity by using techniques well known to those skilled in the art, for example iso-electric focusing, HPLC-based chromatography or affinity chromatography.

Current indications suggest that an epitope can be mimicked by a catamer which is about eight monomers in length when the monomers come from the set of alpha amino acids. The ability of the 8-catamer to be the mimotope of the epitope is not critically dependent on every position having a particular monomer. It is to be understood, however, that the present invention is not restricted to sequences formed from eight monomers.

The plurality of catamer preparations required to be synthesized in the application of this invention may be prepared by any of the known methods of catamer synthesis. The preferred method when the monomers are amino acids is to use the solid phase technique described in Australian Patent Specification No. 25429/84, whereby each mixture of catamers is synthesized on a polyethylene support.

The following is a detailed description of one embodiment of the present invention:-

#### 15 A. Synthesis of a plurality of catamer preparations.

As noted above, the preferred method of applying this invention is to synthesize the catamers on a solid support. In this embodiment, the plurality of catamers will all have the general formula

20  $Y-X-X-D_2-D_1-X-X-X-X$  [solid support],

where 'X' represents a member of a set of monomers. Ideally, each catamer preparation should include catamers in which each of the 'X' positions in the catamer formula represents a member of the set of monomers such that each member of the set is present in the catamer preparation in approximately equimolar amounts. Note that the same set of monomers need not be used at each 'X' position. 'Y' in the general formula is an end group of the catamer and may be, but is not restricted to, for example a hydrogen atom or an acetyl group.  $D_1$  and  $D_2$  represent designated positions occupied by monomers which are known and constant in each catamer preparation synthesized; but which are altered systematically between catamer preparations. Note that the same set of monomers need not be used at each designated position. It should also be noted that when the catamers are synthesized from amino acids the catamers may be coupled to the solid support through their -COOH ends, or through their amine ends.

If  $i$  is the number of members in the set of monomers to be coupled in the  $D_1$  position and  $j$  is the number of members in the set of monomers to be coupled in the  $D_2$  position then a total of  $i \cdot j$  different catamer preparations will be synthesized.

In the present embodiment, the support rods are prepared so that the monomers can be coupled to them. The rods are then exposed to the reaction mixture containing each member of the set of monomers being considered. This will couple monomers at the first 'X' position. This process is repeated thrice to give the catamer  $Z-X-X-X-X$  [solid support] where 'Z' is the protective group appropriate to the set of monomers being considered; examples of 'Z' where the monomers are amino acids include the BOC and Fmoc groups. Up to this stage in the performance of the method all rods have been treated in an identical manner.

For the coupling at the  $D_1$  position each rod will be treated with a reaction mixture which contains only a single monomer such as a protected amino acid or the like. In the  $D_1$  position each of the  $i$  monomers will be coupled to  $j$  rods. For the coupling at the  $D_2$  position each rod is again treated with a reaction mixture containing only a single monomer such as a protected amino acid or the like. Each of the  $j$  rods which has a particular monomer in the  $D_1$  position will have a different monomer coupled at the  $D_2$  position. In this way every combination of the members of the set(s) of monomers will be found in the  $i \cdot j$  rods. Coupling of monomers in the remaining 'X' positions is identical to the coupling in the 'X' positions closer to the solid support.

After synthesis of the plurality of catamer preparations any side-chain protective groups are removed from the catamer preparations using the usual techniques and the rod-coupled catamers are washed.

It has been found to be a preferred embodiment of the invention to synthesize more than one set of pluralities of catamer preparations to aid in the analysis of data. Thus, as well as synthesizing catamers with the general formula

55  $Y-X-X-D_2-D_1-X-X-X-X$  [solid support]

as described above, additional sets of catamers may be prepared with the general formulae

Y-X-X-D<sub>2</sub>-X-D<sub>1</sub>-X-X-X-[solid support]

and

Y-X-D<sub>2</sub>-X-X-D<sub>1</sub>-X-X-X-[solid support].

Note that other general formulae may be selected for the syntheses of other sets of catamer preparations.

#### 10 B. Testing of the catamer preparations.

The plurality of catamer preparations prepared as in A above are then contacted with the particular antibody of interest. The reaction between antibody and components of the catamer preparations can then be detected by any of the usual methods, for example Radioimmunoassay (RIA). However, the preferred  
15 method of detection of the presence of antibody-mimotope binding is to use the well-known enzyme-linked immunosorbent assay (ELISA).

At the end of each assay antibodies can be removed from the catamer preparations by, for example, washing with a solution of 8M urea, 0.1% 2-mercaptoethanol and 0.1% sodium dodecylsulphate followed by several washes in phosphate buffered saline. In this way the plurality of catamer preparations may be used  
20 for testing with many other antibodies.

#### C. Analysis of data.

In the testing of the catamer preparations with antibody it will be found that certain rods will show  
25 detectable binding with the antibody. The catamer preparation coupled to such a reacting rod contains a defined combination of the members of the set(s) of monomers in positions D<sub>1</sub> and D<sub>2</sub>. This combination of monomers is likely to form part of the catamer which makes up a mimotope for the epitope which is complementary to the antibody. Further analysis of the data can be carried out in a number of ways. These include:-

- 30 1. combining and permutating these reacting monomers to create a list of catamers which may include one or more mimotopes for the epitope; each catamer in this list could be regarded as a possible mimotope;
2. taking the most suitable candidates from the list of reacting combinations of monomers (but not necessarily those giving the highest antibody binding activity) and further synthesizing sets of catamer  
35 preparations in which the known reacting combination of monomers is held constant and the monomers at other positions of the catamers are varied systematically; thus in the example above, a suitable set of such catamer preparations would include catamers with the formulae

Y-X-D<sub>3</sub>-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-[solid support]

and

Y-X-X-D<sub>2</sub>-D<sub>1</sub>-D<sub>4</sub>-X-X-X-[solid support],

45 where D<sub>1</sub> and D<sub>2</sub> constitute the reacting combination of monomers and D<sub>3</sub> and D<sub>4</sub> are other designated positions where each of the members of the set of monomers is systematically varied; and

3. combining the results from more than one of the sets of plurality of catamer preparations and decoding to produce a single sequence of monomers, or possibly a very small number of such sequences which may mimic the epitope of interest; this is because, using the embodiment given above,  
50 the reacting combinations of monomers when they are adjacent to each other, the reacting combinations of monomers when they are separated by one position and finally the reacting combinations of monomers which are separated by two positions, are all known; these data may be easily decoded to yield the sequence of the mimotope; this approach will be of greatest benefit when the antibody used to test the plurality of catamer preparations is a monoclonal antibody.

#### 55 D. Synthesis of selected mimotopes.

The selected mimotopes can then be synthesized using similar methods to those described in

Australian Patent Specification No. 25429/84. After the selected catamers have been synthesized they are reacted with the antibody of interest. It is then simple to select the catamer which binds most strongly with the antibody. As a final check that this catamer is the best mimotope of the epitope, the sequence of the catamer may be used as the parent sequence of a replacement net as described in Australian Patent Specification No. 25428/84.

In Examples 1, 2 and 3 given below, the defined set of monomers is the set of the common L-alpha amino acids. This set of monomers is used for both the designated positions and the random positions (positions 'D<sub>1</sub>', 'D<sub>2</sub>' and 'X' as used in the general formulae above). Furthermore, in these examples, the end group of the catamers (the 'Y' group) was the acetyl moiety.

#### EXAMPLE 1

A plurality of 8-catamer preparations was synthesized with the general formula

Acetyl-NH-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-[solid support]

where 'X' is a member of the set of 20 common L-alpha amino acids, such that each member of the set is present at the 'X' position in the catamer preparations in approximately equi-molar amounts; both designated positions, D<sub>1</sub> and D<sub>2</sub> are members of the same set of monomers; and '-NH-' represents the terminal amine group of the catamer. The 8-catamer preparations were synthesized on a solid polymer pin or rod as described in general in Australian Patent Specification No. 25429/84, using the BOC group as protection for the L-alpha amino acids during synthesis.

Members of the set of common L-amino acids were coupled in the 'X' positions in the following way:-

1. A mixture of amino acids was dissolved in 102ml of Dimethylformamide (DMF). The mixture consisted of:-

BOC-L-Alanine	32 mg
BOC-Methoxybenzyl-L-Cysteine	100 mg
BOC-Benzyl-L-Aspartic acid	90 mg
BOC-Benzyl-L-Glutamic acid	96 mg
BOC-L-Phenylalanine	61 mg
BOC-L-Glycine	27 mg
BOC-Tosyl-L-Histidine	142 mg
BOC-L-Isoleucine.1/2 H <sub>2</sub> O	51 mg
BOC-Carbobenzoxy-L-Lysine	124 mg
BOC-L-Leucine H <sub>2</sub> O	55 mg
BOC-L-Methionine	54 mg
BOC-L-Asparagine	47 mg
BOC-L-Proline	42 mg
BOC-L-Glutamine	53 mg
BOC-Tosyl-L-Arginine.H <sub>2</sub> O	167 mg
BOC-Benzyl-L-Serine	76 mg
BOC-Benzyl-L-Threonine	83 mg
BOC-L-Valine	42 mg
BOC-L-Tryptophan	78 mg
BOC-Benzyl-L-Tyrosine	116 mg

2. A solution of 1310 mg of 1-Hydroxybenzotriazole (HOBT) was made in 34 ml of DMF.

3. A solution of 1150 mg of N,N'-Dicyclohexylcarbodiimide (DCC) was made in 34 ml of DMF.

4. Before use, the solution of HOBT was added to the solution of amino acids and mixed. Then the solution of DCC was added and mixed. The catamer or linking molecule already on the solid support was then reacted with the resulting mixture.

Individual L-amino acids were coupled in the D<sub>1</sub> and D<sub>2</sub> positions by the method described in Australian Patent Specification No. 25429/84. This resulted in the synthesis of 400 catamer preparations, comprising all possible pairings of the 20 common amino acids at the D<sub>1</sub> and D<sub>2</sub> positions.

The plurality of 8-catamer preparations were tested against a monoclonal antibody by ELISA. This monoclonal antibody was prepared against Foot-and-Mouth Disease Virus (FMDV) subtype A<sub>10</sub> using

techniques well known to those skilled in the art. In performing the ELISA assay support-coupled peptides were incubated in a precoat solution of 1% ovalbumin/1% bovine serum albumin/0.1% Tween 20 in phosphate buffered saline (PBS) for 1 hr at 25°C to block non-specific absorption of antibodies. Overnight incubation at 4°C in a suitable dilution of monoclonal antibody in precoat solution was followed by three washes in 0.05% Tween 20/PBS. Reaction for 1 hr at 25°C with goat anti-mouse IgG conjugated to horseradish peroxidase, diluted in the precoat solution, was again followed by extensive washing with PBS/Tween to remove excess conjugate. The presence of antibody was detected by reaction for 45 min with a freshly prepared enzyme substrate solution (40mg of o-phenylenediamine and 18μl of "120 vol." hydrogen peroxide in 100ml of phosphate/citrate buffer, pH 5.0), and the colour produced determined at 450nm.

Figure 1 shows the result of testing the 8-catamer preparations with this antibody. In Figure 1, each group of twenty lines represents the response in an ELISA in units of absorbancy of the colour developed when measured at 450nm. Each member of the group of twenty lines has a common amino acid in the 'D<sub>1</sub>' position which is given in single-letter code immediately below each group. Each member of a group of twenty lines represents a different amino acid in the 'D<sub>2</sub>' position - these are in alphabetical order of their single-letter code. The results are summarised in Table 1. (In Table 1, and throughout the remainder of the Examples, the single letter code for amino acids will be used.)

TABLE 1

Antibody binding activities of the 8-catamer preparations mixtures Ac-X-X-D <sub>2</sub> -D <sub>1</sub> -X-X-X-X-S <sub>s</sub> .	
Peptide Mixture	Extinction E <sub>415</sub>
Ac-X-X-M-H-X-X-X-X-S <sub>s</sub>	0.624
Ac-X-X-M-K-X-X-X-X-S <sub>s</sub>	0.582
Ac-X-X-K-K-X-X-X-X-S <sub>s</sub>	0.537
Ac-X-X-K-H-X-X-X-X-S <sub>s</sub>	0.510
Ac-X-X-Q-H-X-X-X-X-S <sub>s</sub>	0.488
Ac-X-X-H-H-X-X-X-X-S <sub>s</sub>	0.486
Ac-X-X-Q-N-X-X-X-X-S <sub>s</sub>	0.460
Ac-X-X-W-M-X-X-X-X-S <sub>s</sub>	0.448
Ac-X-X-K-N-X-X-X-X-S <sub>s</sub>	0.431
Ac-X-X-H-K-X-X-X-X-S <sub>s</sub>	0.417
Mean extinction of the lower 200 values was 0.202 (Standard deviation 0.021). Ac means the acetyl moiety. S <sub>s</sub> means [solid support].	

It will be seen that the antibody reacts strongly with a number of 8-catamer preparations. In order of strength of reaction, these include 8-catamer preparations in which the designated positions (-D<sub>2</sub>-D<sub>1</sub>-) are: -M-H-, -M-K-, -K-K-, -K-H-, -Q-H- and -H-H-.

Two further pluralities of 8-catamer preparations were synthesized. These had the formulae

Acetyl-NH-X-D<sub>3</sub>-M-K-X-X-X-X-[solid support]

and

Acetyl-NH-X-X-M-K-D<sub>4</sub>-X-X-X-X-[solid support]

where 'M' means a methionine residue, 'K' means a lysine residue and 'D<sub>3</sub>' and 'D<sub>4</sub>' are the new designated position for these series of syntheses; other terms in the formulae are as used previously.

These 8-catamer preparations were tested against the monoclonal antibody against FMDV A<sub>10</sub>. Figures 2A and 2B show the results of ELISA testing with the monoclonal antibody used to produce the results shown in Figure 1. Figure 2A represents 8-catamer preparations with the general formula

Acetyl-NH-X-D<sub>3</sub>-M-K-X-X-X-X-[solid support].

Figure 2B represents 8-catamer preparations with the general formula

Acetyl-NH-X-X-M-K-D<sub>3</sub>-X-X-X-[solid support].

- 5 In both figures the amino acid in the '-D<sub>3</sub>-' or '-D<sub>4</sub>-' position is given below each line which represents the absorbancy of the colour developed when measured at 450nm. The results are summarised in Table 2. Antibody binding activities are shown relative to the 8-catamer preparations containing the defined amino acid pair M-K for all the 8-catamer preparations giving a value greater than 1.1. Also shown is the value found for the reverse pair K-M.

10  
15  
TABLE 2 Relative antibody binding activities of 8-catamer preparations containing three defined amino acids.

Amino-terminal extension		Carboxyl-terminal extension	
Sequence	Activity	Sequence	Activity
Ac-X-H-M-K-X-X-X-X-S <sub>s</sub>	2.54	Ac-X-X-M-K-H-X-X-X-S <sub>s</sub>	2.85
Ac-X-W-M-K-X-X-X-X-S <sub>s</sub>	2.50	Ac-X-X-M-K-Q-X-X-X-S <sub>s</sub>	2.20
Ac-X-Q-M-K-X-X-X-X-S <sub>s</sub>	1.69	Ac-X-X-M-K-W-X-X-X-S <sub>s</sub>	1.97
Ac-X-N-M-K-X-X-X-X-S <sub>s</sub>	1.27	Ac-X-X-M-K-G-X-X-X-S <sub>s</sub>	1.41
Ac-X-G-M-K-X-X-X-X-S <sub>s</sub>	1.14	Ac-X-X-M-K-N-X-X-X-S <sub>s</sub>	1.26
Ac-X-C-M-K-X-X-X-X-S <sub>s</sub>	1.11		
Control sequences M-K		1.00 (E <sub>450</sub> = 0.14)	
K-M		0.97	
Test background E <sub>450</sub> = 0.08.			

40 It will be seen that the 8-catamer preparations which gave the greatest reactions with the monoclonal antibody had the following sequences in the designated positions: -H-M-K-, -W-M-K-, and -Q-M-K- in Figure 2A; and -M-K-H-, -M-K-Q and -M-K-W- in Figure 2B.

45 Incorporation of histidine at either side of the M-K defined pair resulted in the greatest antibody binding activity. However histidine at the amino terminal side was not significantly better than tryptophan, and the defined sequence W-M-K-H was selected for further evaluation.

The validity of proceeding with the defined sequence W-M-K-H, predicted to be optimal by combining the individual preferred extensions to the pair M-K, was tested. Catamer preparations based on the parent sequence of Acetyl-NH-X-W-M-K-H-X-X-X-S<sub>s</sub> [solid support] were synthesized, in which at each of the four defined positions all 19 alternative amino acids were substituted, one at a time. Testing of this set of catamer preparations for binding activity with the test monoclonal antibody (MAb) showed Q-M-R-H to be the preferred sequence for extension (E<sub>450</sub> obtained were; Ac-X-W-M-K-H-X-X-X-S<sub>s</sub> = 0.69, Ac-X-Q-M-K-H-X-X-X-S<sub>s</sub> = 1.47, and Ac-X-W-M-R-H-X-X-X-S<sub>s</sub> = 1.11, test background was 0.2). The preference for glutamine at the amino-terminal side of the defined sequence M-K-H rather than tryptophan as had been 55 determined for the pair M-K, demonstrates the inter-dependence of individually determined optima.

8-Catamer preparations with the general formulae

Acetyl-NH-D<sub>5</sub>-Q-M-R-H-X-X-X- [solid support] and



Acetyl-NH-X-Q-M-R-H-D<sub>5</sub>-X-X- [solid support]

were synthesized. The preparations with the highest antibody-binding activity were those with D<sub>5</sub> = tryptophan and D<sub>6</sub> = serine, which suggests that W-Q-M-R-H-S is the optimum sequence. No significant improvement in antibody binding activity was achieved by defining further amino acids to either end of this sequence. Assessment of the set of peptides comprising all the single amino acid substitutions of this sequence, synthesized as hexapeptides, confirmed that W-Q-M-R-H-S was close to the optimum hexapeptide. Some increase in antibody binding activity was shown when glycine replaced arginine in the peptide to give the sequence W-Q-M-G-H-S.

In order to confirm the specificity of the antibody-binding reaction of the determined group of related peptides for the test MAb, the titre of the test MAb for a number of the peptides was compared with those of a MAb to Sperm whale myoglobin or hepatitis A (Table 3). The specificity of the antibody-binding reaction is clearly for the test MAb, the antibody used to determine the sequence of those peptides.

TABLE 3

Specificity of Test MAb for W-Q-M-G-H-S related peptides.			
Peptide	Titre		
	Test MAb	anti-hepatitis A	anti-myoglobin
Ac-X-X-M-K-X-X-S <sub>s</sub>	2,000	< 200	< 200
Ac-X-W-M-K-X-X-S <sub>s</sub>	14,000	< 200	1,600
Ac-X-X-M-K-H-X-S <sub>s</sub>	17,000	< 200	< 200
Ac-X-W-M-K-H-X-S <sub>s</sub>	36,000	< 200	800
Ac-W-Q-M-R-H-S-S <sub>s</sub>	100,000	< 200	< 200
Ac-W-Q-M-G-H-S-S <sub>s</sub>	120,000	< 200	< 200

Antibody titres for each MAb were determined by ELISA and correspond to the reciprocal dilution of the ascites fluid giving an extinction of three times the test background. Values were corrected for non-specific absorption as determined by reacting each dilution of ascites fluid with unrelated peptide controls (Ac-G-D-L-G-S-I-S<sub>s</sub> and Ac-G-D-L-Q-V-L-S<sub>s</sub>). Monoclonal antibodies tested were: Test MAb, anti-FMDV, type A<sub>10</sub> as described in the text (titre against the whole virus 1.3x10<sup>5</sup>); anti-hepatitis A, site of reaction unknown (titre against hepatitis A virus 10<sup>6</sup>); and anti-myoglobin (sperm whale), site of reaction also unknown (titre against sperm whale myoglobin 10<sup>6</sup>).

## EXAMPLE 2

Example 2 illustrates the application of the invention to a polyclonal antibody.  
A further plurality of 8-catamers was synthesized with the general formula

Acetyl-NH-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-[solid support]

as described in Example 1. This was tested against an antibody which had been produced against the synthetic peptide C-G-D-L-G-S-I-A-K. This peptide was prepared using the usual techniques of Merrifield solid-phase peptide synthesis. The peptide was coupled to Keyhole Limpet Haemocyanin through the sulfhydryl group of the cysteine moiety. The coupled peptide was formulated with Freund's incomplete adjuvant and injected into a rabbit via the intramuscular route. Antiserum was obtained by bleeding the rabbit 40 days after injection.

The results of ELISA testing the plurality of 8-catamer preparations with the antibody are given in Figure 3 in the same manner as in Figure 1. It will be seen that all 8-catamer preparations which had E (glutamic acid) in the D<sub>1</sub> position reacted significantly with antibody. Other combinations which reacted significantly with antibody include in decreasing order:- -D-I-, -G-D-, -I-G-, and -D-V-.

As described in Example 1 above, further pluralities of 8-catamer preparations were synthesized with the formulae:-

Acetyl-NH-X-D<sub>3</sub>-D-I-X-X-X-X-[solid support]

and

Acetyl-NH-X-X-D-I-D<sub>4</sub>-X-X-X-[solid support]

- 5 where D is aspartic acid and I is isoleucine. Results of testing with the antibody are shown in Figure 4 in the same manner as in Figure 2. Figure 4A represents 8-catamer preparations with the general formula

Acetyl-NH-X-D<sub>3</sub>-D-I-X-X-X-[solid support].

- 10 Figure 4B represents 8-catamer preparations with the general formula

Acetyl-NH-X-X-D-I-D<sub>4</sub>-X-X-X-[solid support].

- Figure 4 shows that 8-catamer preparations which react with antibody include the following sequences  
15 in the designated positions:- -G-D-I-, -A-D-I-, -P-D-I-, -D-I-D-, -D-I-M- and -D-I-T-. It can plainly be seen by comparing Figure 4A with Figure 4B that the position immediately before the -D-I- pair permits far less latitude in allowable amino acids than does the position immediately after the -D-I- pair.

8-Catamer preparations based on the parent sequence

- 20 Acetyl-NH-X-X-G-D-I-D-X-X-[solid support]

were synthesized, in which at each of the four defined positions in the catamer preparation all 19 alternative amino acids were substituted, one at a time. Testing of this set of preparations for binding activity with the test antibody showed that to retain the ability to react with this antibody, the glycine residue could only be  
25 replaced with alanine; the amino-terminal aspartic acid could not be replaced by any amino acid; the isoleucine could be replaced with valine, leucine or threonine or with about ten other amino acids with a reduction of binding activity; and the carboxyl-terminal aspartic acid could be replaced by virtually any amino acid, serine, in this position giving the strongest binding. The testing showed that the defined sequence G-D-I-S was the preferred one for extension.

- 30 Extending the defined sequence above demonstrated the preference for incorporation of tryptophan at the amino-terminal end and histidine at the carboxyl-terminal end. However, neither extension produced a very marked increase in the antibody-binding activity.

Further extension of the preferred defined sequence W-G-D-I-S-H showed that there was little advantage in extending the sequence further.

- 35 It will be noted that the mimotope determined for this particular antibody, namely, W-G-D-I-S-H has only a passing similarity to the peptide which was used to induce the antibody, namely, C-G-D-L-G-S-I-A-K. However, testing of the antibody in a replacement net based on the immunogenic sequence G-D-L-G-S-I showed a marked preference for the sequence G-D-L-G-D-I rather than the inducing sequence G-D-L-G-S-I (E<sub>450</sub> values 0.45 and 0.16 respectively). There is a similarity between the mimotope as determined and a  
40 preferred binding peptide for the antibody. This is strong evidence for the broad applicability of the method.

### EXAMPLE 3

- Example 3 illustrates that the length of the plurality of catamer preparations can be shorter than 8-  
45 catamers.

A plurality of 4-catamer preparations was synthesized with the general formula

Acetyl-NH-X-D<sub>2</sub>-D<sub>1</sub>-X-[solid support]

- 50 where the abbreviations are as used for Example 1. The method of synthesis was identical to that used for Example 1. This was tested against a monoclonal antibody which had been raised against sperm whale myoglobin using techniques well known to those skilled in the art.

The results of ELISA testing the plurality of 4-catamer preparations with the antibody are given in Figure 5 in the same manner as in Figure 1. It will be seen that only the catamer preparations with the formula

- 55 Acetyl-NH-X-L-A-X-[solid support]

was able to react with the monoclonal antibody.

As described in Example 1 above, further pluralities of 4-catamer preparations were synthesized with the formulae:

Acetyl-NH-D<sub>3</sub>-L-A-X-[solid support]

and

Acetyl-NH-X-L-A-D<sub>4</sub>-[solid support]

Table 4 summarizes the most reactive catamer preparations when tested in an ELISA system with the monoclonal antibody.

**TABLE 4 - Relative antibody binding activities of 4-catamer preparations containing three defined amino acids.**

Amino-terminal extension		Carboxyl-terminal extension	
Sequence	Activity	Sequence	Activity
AC-P-L-A-X-S <sub>5</sub>	1.80	Ac-X-L-A-Q-S <sub>5</sub>	1.73
Ac-A-L-A-X-S <sub>5</sub>	1.36	Ac-X-L-A-R-S <sub>5</sub>	1.58
Ac-F-L-A-X-S <sub>5</sub>	1.33	Ac-X-L-A-S-S <sub>5</sub>	1.39
Control sequences Ac-X-L-A-X-S <sub>5</sub> 1.00 (E <sub>450</sub> = 0.17)			
Ac-X-A-L-X-S <sub>5</sub> 0.94			

Catamer preparations based on the parent sequence

Acetyl-NH-P-L-A-Q-[solid support]

were synthesized to form a replacement net as described in Example 2. Figure 6 shows the result of testing these peptides in an ELISA with the monoclonal antibody. Each group of twenty lines in Figure 6 represents the response in percentage when compared to the mean of the parent response (E<sub>450</sub> = 2.62 with a background of 0.25). Each member of a group of twenty lines has had the amino acid indicated below substituted by another amino acid. The order of the lines is the alphabetical order of the single letter code of the amino acid. The thickened line in each group of twenty indicates the parent sequence P-L-A-Q. It can clearly be seen that for the catamer preparation to be able to bind to this monoclonal antibody with at least half the activity of the parent sequence the proline may be replaced with the valine, isoleucine, leucine, alanine or serine; the leucine can be replaced by phenylalanine only; the alanine is freely replaceable by virtually any other amino acid; and the glutamine residue can only be replaced by asparagine.

A set of all overlapping 4-catamers of the sperm whale myoglobin sequence was synthesized using the methods described for the designated positions in Example 1. These catamers were reacted in an ELISA with the monoclonal antibody. The results are shown in Figure 7. Each line represents the colour developed in an ELISA, read at 450nm. Each line is numbered so that the residue at the amino terminal end of the particular catamer has the same number as in the sperm whale myoglobin sequence.

It can clearly be seen that the only 4-catamer which reacts with the monoclonal antibody is that comprising residues 88 to 91 of the sperm whale sequence. This sequence is P-L-A-Q. Clearly, the monoclonal antibody recognizes this particular determinant on the sperm whale myoglobin.

A set of 5-catamers with the general formulae

Acetyl-NH-D<sub>5</sub>-P-L-A-Q-[solid support]

5 and

Acetyl-NH-P-L-A-Q-D<sub>5</sub>-[solid support]

10 were synthesized to determine the best extensions to the defined sequence. The best binding preparations were

Acetyl-NH-K-P-L-A-Q-[solid support]  
and Acetyl-NH-P-L-A-Q-Y-[solid support].

15 It should be noted that the sequence K-P-L-A-Q is homologous with the sequence of sperm whale myoglobin at residues 87 to 91. Again, this example illustrates the applicability of this technique.

### Claims

- 20 1. A method of determining the sequence of monomers selected from a defined set of monomers which is a conformational equivalent of an epitope which is complementary to a particular paratope of an antibody of interest, the method comprising the steps of:
  - (i) synthesizing a plurality of catamer preparations; each of said catamer preparations consisting of a plurality of catamers made of a defined set of monomers and in which (i) the composition at two or more designated positions in each catamer is known and is constant for each catamer in the catamer preparation, and (ii) the composition at each of the remaining positions are randomly made up from the members of said defined set of monomers; and said plurality of catamer preparations comprising preparations in which the monomer(s) at the designated positions are systematically varied to contain members from said defined set of monomers;
  - 25 (ii) contacting each of said plurality of catamer preparations with the antibody of interest; and
  - (iii) detecting the presence or absence of binding between each of said plurality of catamer preparations and said given antibody; and
  - (iv) determining a partial sequence of monomers for the conformational equivalent of an epitope which is complementary to the particular paratope of said antibody from the known composition of said two or more designated positions in each of said plurality of catamer preparations that bind to said antibody.
- 30 2. A method according to claim 1, further comprising the additional steps of:
  - (v) synthesizing a further plurality of catamer preparations as defined in claim 1 and which include the composition of the partial sequence as determined in said step (iv) as initial designated position, each of said further plurality of catamer preparations having at least one or more additional position in each catamer as a designated position in which the monomer is known; and
  - (vi) repeating (ii), (iii), and (iv) as defined in claim 1 in respect of said further plurality of catamer preparations.
- 45 3. A method according to claim 1, wherein each of said plurality of catamer preparations is made up from the set of alpha-amino acids.
4. A method according to claim 3, wherein each of said plurality of catamer preparations consists of a plurality of 4-catamers to 8-catamers.
- 50 5. A method according to claim 1, wherein each of said plurality of catamers is synthesized on a solid support, and has the general formula:
 

55 Y-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-[solid support],  
Y-X-X-D<sub>2</sub>-X-D<sub>1</sub>-X-X-X-[solid support]  
or Y-X-D<sub>2</sub>-X-X-D<sub>1</sub>-X-X-X-[solid support]

wherein Y is an end group of the catamer, X (which may be the same or different) is a member of the set of monomers such that each member is present in the catamer preparation, and D<sub>1</sub> and D<sub>2</sub> are known monomers which are constant in each of the catamer preparations.

- 5 6. A method according to claim 5, wherein each of said plurality of catamers has the general formula:

Acetyl-NH-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-[solid support]

10 wherein X is a member of the set of L-alpha amino acids, and such plurality of catamers is such that each member of the set is present at the X position in approximately equi-molar amounts, D<sub>1</sub> and D<sub>2</sub> are known monomers selected from the set of L-alpha amino acids.

### Patentansprüche

- 15 1. Verfahren zur Sequenzbestimmung von Monomeren, ausgewählt aus einem definierten Satz von Monomeren, welcher ein Konformations-Äquivalent eines Epitops ist, das komplementär zu einem bestimmten Paratop eines Antikörpers von Interesse ist, wobei das Verfahren die Schritte umfasst:
- (i) eine Vielzahl von Catamer-Präparationen herzustellen; wobei jede der Catamer-Präparationen aus einer Vielzahl von Catameren besteht, die aus einem definierten Satz von Monomeren hergestellt wurden und worin (i) die Zusammensetzung an zwei oder mehr bezeichneten Positionen in jedem Catamer bekannt und für jedes Catamer in der Catamer-Präparation konstant ist, und (ii) die Zusammensetzung der verbleibenden Positionen durch die Teile des definierten Satzes der Monomere zufällig gebildet werden; und die Vielzahl der Catamer-Präparationen umfassen, in denen das/die Monomer(e) bei den bezeichneten Positionen systematisch variiert werden, damit Teile des definierten Satzes von Monomeren enthalten sind;
- 20 (ii) Zusammenbringen jeder der Vielzahl von Catamer-Präparationen mit einem Antikörper von Interesse; und
- (iii) Nachweis des Vorhandenseins oder Fehlens von Bindung zwischen jeder der Vielzahl von Catamer-Präparationen und dem gegebenen Antikörper; und
- 25 (iv) Bestimmung einer Teil-Sequenz von Monomeren aus der bekannten Zusammensetzung der zwei oder mehr bezeichneten Positionen in jeder der Vielzahl der an den Antikörper bindenden Catamer-Präparationen, für das Konformations-Äquivalent eines Epitops, das zu dem einzelnen Paratop des Antikörpers komplementär ist.
- 30 2. Verfahren nach Anspruch 1, weiter umfassend die zusätzlichen Schritte:
- (v) eine weitere Vielzahl von Catameren wie in Anspruch 1 definiert, und welche die Zusammensetzung der Teilsequenz wie in Schritt (iv) als anfänglich bezeichnete Positionen bestimmt, einschließlich, herzustellen, wobei jede weitere Vielzahl von Catamer-Präparationen wenigstens eine oder mehrere zusätzliche Positionen in jedem Catamer als eine bezeichnete Position, an der das Monomer bekannt ist, enthält; und
- 35 (vi) (ii), (iii), und (iv) wie in Anspruch 1 beschrieben zu wiederholen, im Hinblick auf die weitere Vielzahl von Catamer-Präparationen.
3. Verfahren nach Anspruch 1, wobei jede der Vielzahl der Catamer-Präparationen aus dem Satz der alpha-Aminosäuren gebildet wird.
- 40 4. Verfahren nach Anspruch 3, wobei jede der Vielzahl der Catamer-Präparationen aus einer Vielzahl von 4-Catameren bis 8-Catameren besteht.
5. Verfahren nach Anspruch 1, wobei jede Vielzahl von Catameren an einem festen Träger hergestellt wird, und die allgemeine Formel besitzt:
- Y-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-(fester Träger)
- Y-X-X-D<sub>2</sub>-X-D<sub>1</sub>-X-X-X-(fester Träger)
- 55 Y-X-D<sub>2</sub>-X-X-D<sub>1</sub>-X-X-X-(fester Träger)

wobei y eine endständige Gruppe des Catamers ist, X (welches gleich oder unterschiedlich sein kann) ein Teil des Satzes von Monomeren ist, dergestalt, daß jeder Teil in der Catamer-Präparation

vorhanden ist, und D<sub>1</sub> und D<sub>2</sub> bekannte Monomere sind, die in jeder Catamer-Präparation konstant sind.

6. Verfahren nach Anspruch 5, wobei jede Vielzahl von Catameren die allgemeine Formel besitzt:

Acetyl-NH-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-(fester Träger)

wobei X ein Teil des Satzes von L-alpha-Aminosäuren ist, und diese Vielzahl der Catamere dergestalt ist, daß jeder Teil des Satzes an der X-Position in etwa äquimolaren Mengen vorhanden ist, D<sub>1</sub> und D<sub>2</sub> bekannte Monomere ausgewählt aus dem Satz von L-alpha-Aminosäuren, sind.

## Revendications

1. Procédé de détermination de la séquence de monomères choisis dans un ensemble défini de monomères qui est un équivalent conformationnel d'un épitope complémentaire d'un paratope particulier d'un anticorps intéressant, le procédé comprenant les étapes de :

(i) synthèse d'une pluralité de préparations de catamères; chacune desdites préparations de catamères consistant en une pluralité de catamères constituée d'un ensemble défini de monomères et dans laquelle (i) la composition en ce qui concerne deux positions désignées ou plus dans chaque catamère est connue et constante pour chaque catamère dans la préparation de catamères, et (ii) la composition sur chacune des positions restantes est constituée de façon aléatoire des membres dudit ensemble défini de monomères; et ladite pluralité de préparations de catamères comprend des préparations dans lesquelles le ou les monomères aux positions désignées sont systématiquement modifiés pour contenir des membres dudit ensemble défini de monomères;

(ii) mise en contact de chacune de ladite pluralité de préparations de catamères avec l'anticorps intéressant; et

(iii) détection de la présence ou de l'absence de liaison entre chacune de ladite pluralité de préparations de catamères et ledit anticorps donné; et

(iv) détermination d'une séquence partielle de monomères pour l'équivalent conformationnel d'un épitope complémentaire du paratope particulier dudit anticorps à partir de la composition connue desdites deux positions désignées ou plus dans chacune de ladite pluralité de préparations de catamères liée audit anticorps.

2. Procédé selon la revendication 1, comprenant de plus les étapes supplémentaires de :

(v) synthèse d'une pluralité supplémentaire de préparations de catamères telle que définie dans la revendication 1 et qui comprend la composition de la séquence partielle telle que déterminée dans ladite étape (iv) comme position désignée initiale, chacune de ladite pluralité supplémentaire de préparations de catamères ayant au moins une position additionnelle ou plus dans chaque catamère comme position désignée dans laquelle le monomère est connu; et

(vi) la répétition de (ii), (iii), et (iv) telles que définies dans la revendication 1 en ce qui concerne ladite pluralité supplémentaire de préparations de catamères.

3. Procédé selon la revendication 1, dans lequel chacune de ladite pluralité de préparations de catamères est constituée de l'ensemble des alpha-aminoacides.

4. Procédé selon la revendication 3, dans lequel chacune de ladite pluralité de préparations de catamères consiste en une pluralité de 4-catamères à 8-catamères.

5. Procédé selon la revendication 1, dans lequel chacune de ladite pluralité de catamères est synthétisée sur un support solide, et a la formule générale :

Y-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-X-[support solide],

Y-X-X-D<sub>2</sub>-X-D<sub>1</sub>-X-X-X-[support solide]

ou Y-X-D<sub>2</sub>-X-X-D<sub>1</sub>-X-X-X-[support solide]

dans laquelle Y est un groupe terminal du catamère, X (qui peut être identique ou différent) est un membre de l'ensemble de monomères tel que chaque membre est présent dans la préparation de catamères, et D<sub>1</sub> et D<sub>2</sub> sont des monomères connus qui sont constants dans chacune des préparations.


de catamères.

6. Procédé selon la revendication 5, dans lequel chacune de ladite pluralité de catamères a la formule générale :

Acétyl-NH-X-X-D<sub>2</sub>-D<sub>1</sub>-X-X-X-[support solide]


dans laquelle X est un membre de l'ensemble des L-alpha aminoacides, cette pluralité de catamères étant telle que chaque membre de l'ensemble est présent à la position X en quantités approximativement équimolaires, D<sub>1</sub> et D<sub>2</sub> étant des monomères connus choisis dans l'ensemble des L-alpha aminoacides.

FIGURE 1






# FIGURE 2



A



B

FIGURE 3




FIGURE 4








FIGURE 5



## FIGURE 6



# FIGURE 7



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**